A Reliable Research Partner in Life Science and Medicine

Recombinant Human CSF2RA/GM-CSFR Protein (His Tag)

Catalog Number: PKSH033281

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip					
	00	CI	411	n	П	n	m
JU	\mathbf{c}	U		J A	ш	v.	ш

Species Human

Source HEK293 Cells-derived Human CSF2RA/GM-CSFR protein Glu23-Gly320, with an C-

terminal His

Calculated MW35.5 kDaObserved MW60 kDaAccessionP15509

Bio-activity Measured by its ability to inhibit GM-CSF-dependent proliferation of TF- 1 human

erythroleukemic cells. The ED_{50} for this effect is 0.5-2 µg/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Granulocyte-Macrophage Colony-Stimulating Factor Receptor Subunit α (CSF2RA) is a single-pass type I membrane protein which belongs to the type I cytokine receptor family of Type 5 subfamily. The CSF2RA gene is found in the pseudoautosomal region (PAR) of the X and Y chromosomes with some of the isoforms being membrane-bound and others being soluble. CSF2RA is a low affinity receptor for granulocyte-macrophage colony-stimulating factor. CSF2RA transduces a signal that results in the proliferation, differentiation, and functional activation of hematopoietic cells. Defects in CSF2RA are the cause of pulmonary surfactant metabolism dysfunction type 4 (SMDP4).

For Research Use Only