## Recombinant Human ICOS/AILIM Protein (Fc Tag)

Catalog Number: PKSH033644



Description		
Species	Human	
Mol_Mass	40.9 kDa	
Accession	Q9Y6W8	
<b>Bio-activity</b>	Not validated for activity	
Properties		
Purity	>95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa	МК	R
120 90 60 40		-
30	-	
20	-	
14	-	

> 95 % as determined by reducing SDS-PAGE.

## Background

Inducible T-cell costimulator, also known as activation-inducible lymphocyte immunomediatory molecule, CD278, AILI M, CVID1 and ICOS, belongs to the CD28 and CTLA4 cell surface receptor family.. ICOS contains one Ig-like V-type domain and exsits as a homodimer with disulfide-linked. ICOS is highly expressed on tonsillar T-cellsand can be induced by PMA and ionomycin, ICOS plays an important role in cell-cell signaling, immune responses, and regulation of cell proliferation. Defects in ICOS are the cause of immunodeficiency common variable type 1, which is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antige

## For Research Use Only