
Caspase-3 Polyclonal Antibody

Catalog Number:E-AB-63510 1 Publications




Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human,Mouse,Rat
Immunogen	A synthetic peptide of human Caspase-3 (NP_004337.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applications	Recommended Dilution
IHC	1:50-1:200
IF	1:50-1:200
Data	




Immunohistochemistry of paraffin-embedded Human lung cancer using Caspase-3 Polyclonal Antibody at dilution of 1:100 (40x lens).




heart using Caspase-3 Polyclonal Antibody at dilution of 1:100 (40x lens).




Immunohistochemistry of paraffin-embedded Human gastric cancer using Caspase-3 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Rat testis using Caspase-3 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Rat kidney using Caspase-3 Polyclonal Antibody at



Immunohistochemistry of paraffin-embedded Rat spleen using Caspase-3 Polyclonal Antibody at

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Caspase-3 Polyclonal Antibody

Catalog Number:E-AB-63510 1 Publications



dilution of 1:100 (40x lens).

dilution of 1:100 (40x lens).

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein.

For Research Use Only

A Reliable Research Partner in Life Science and MedicineToll-free: 1-888-852-8623Tel: 1-832-243-6086Web: www.elabscience.comEmail: techsupport@elabscience.com