

## Recombinant Human FLT3LG/Flt3 Ligand Protein (His Tag)

**Catalog Number: PKSH032452**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human FLT3LG;Flt3 Ligand protein Thr 27-Pro184, with an C-terminal His
<b>Calculated MW</b>	19.0 kDa
<b>Observed MW</b>	24-32 kDa
<b>Accession</b>	P49771
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 0.01 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Fms-Related Tyrosine Kinase 3 Ligand (FLT3LG) is a hematopoietic four helical bundle cytokine. Mature human Flt-3 Ligand consists of an extracellular domain (ECD) with a cytokine-like domain and a juxtamembrane tether region, a transmembrane segment, and a cytoplasmic tail. Human and mouse Flt-3 Ligand show cross-species activity. Flt-3 Ligand is expressed as a noncovalently-linked dimer by T cells and bone marrow and thymic fibroblasts. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors by activation of Flt 3 receptor.

### For Research Use Only