

Recombinant Human FLT3LG/Flt3 Ligand Protein (His Tag)

Catalog Number: PKSH032452

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human FLT3LG;Flt3 Ligand protein Thr 27-Pro184, with an C-terminal His
Mol_Mass	19.0 kDa
Accession	P49771
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.01 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Fms-Related Tyrosine Kinase 3 Ligand (FLT3LG) is a hematopoietic four helical bundle cytokine. Mature human Flt-3 Ligand consists of an extracellular domain (ECD) with a cytokine-like domain and a juxtamembrane tether region, a transmembrane segment, and a cytoplasmic tail. Human and mouse Flt-3 Ligand show cross-species activity. Flt-3 Ligand is expressed as a noncovalently-linked dimer by T cells and bone marrow and thymic fibroblasts. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). In synergy with other growth factors, Flt3 ligand stimulates the proliferation and differentiation of various blood cell progenitors by activation of Flt 3 receptor.

For Research Use Only