

# Recombinant Human Thioredoxin/TXN Protein (His Tag)

Catalog Number: PKSH033108



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Human
<b>Mol_Mass</b>	13.9 kDa
<b>Accession</b>	P10599
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 1mM EDTA, 2mM DTT, pH 7.2. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Thioredoxin (TXN) is a member of the Thioredoxin family. Thioredoxin exists as a disulfide-linked homodimer and contains one Thioredoxin domain. Thioredoxin is up-regulated by ionizing radiation. Thioredoxin participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. Thioredoxin also plays a role in the reversible S-nitrosylation of cysteine residues in target proteins; and thereby contributes to the response to intracellular nitric oxide.

## For Research Use Only