Recombinant Human CD6 Protein(His Tag)

Catalog Number: PDMH100244

Note: Centrifuge before opening to ensure complete recovery of vial contents.

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Description	
Calculated MW42.2 kDaObserved MW60-80 kDaAccessionP30203Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Species	Human
Observed MW60-80 kDaAccessionP30203Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Source	Mammalian-derived Human CD6 protein His 18-Leu402, with an C-terminal His
AccessionP30203Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Calculated MW	42.2 kDa
Bio-activityNot validated for activityPropertiesPurity> 90% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Observed MW	60-80 kDa
Properties Purity > 90% as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -8 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Accession	P30203
Purity> 90% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -8 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Bio-activity	Not validated for activity
Endotoxin< 1.0 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -8 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Properties	
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -8 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	> 90% as determined by reducing SDS-PAGE.
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Shippingreconstituted samples are stable at < -20°C for 3 months.	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.
Mannitol.	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
	Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%
Description for the second state of the second		Mannitol.
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
0.5 mg/mL. Concentration is measured by UV-Vis.		0.5 mg/mL. Concentration is measured by UV-Vis.

Data

kDa	м	R	
80 60	11		
40	-		
30			
20			

SDS-PAGE analysis of Human CD6 proteins, 2 µg/lane of Recombinant Human CD6 proteins was resolved with an SDS-PAGE under reducing conditions, showing bands at 42.2KD

Background

This gene encodes a protein found on the outer membrane of T-lymphocytes as well as some other immune cells. The encoded protein contains three scavenger receptor cysteine-rich (SRCR) domains and a binding site for an activated leukocyte cell adhesion molecule. The gene product is important for continuation of T cell activation. This gene may be associated with an susceptibility to multiple sclerosis (PMID: 19525953, 21849685). Multiple transcript variants encoding different isoforms have been found for this gene.

For Research Use Only Toll-free: 1-888-852-8623

Web:www.elabscience.com