

# Recombinant Human Tachykinin-3/TAC3 Protein (His Tag)

Catalog Number: PKSH033098



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Species</b>	Human
<b>Mol_Mass</b>	13.9 kDa
<b>Accession</b>	Q9UHF0
<b>Bio-activity</b>	Not validated for activity

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Tachykinin 3 (TAC3) is a secreted protein that belongs to the Tachykinin family. Tachykinins are active peptides that excite neurons and evoke behavioral responses; they are potent vasodilators and secretagogues, and contract many smooth muscles in pregnancy. TAC3 is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. It is also expressed in the outer syncytiotrophoblast of the placenta and may be associated with pregnancy-induced hypertension and pre-eclampsia. TAC3 acts as the ligand for the neurokinin-3 receptor, mutations in this gene are associated with normosmic hypogonadotropic hypogonadism.

## For Research Use Only