

Recombinant Human Tetherin/BST2 Protein (His Tag)

Catalog Number: PKSH033296

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|---|
| Species | Human |
| Source | HEK293 Cells-derived Human Tetherin/BST2 protein Asn49-Ser161, with an C-terminal His |
| Calculated MW | 13.7 kDa |
| Observed MW | 23 kDa |
| Accession | Q10589 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. |
| | Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Bone Marrow Stromal Antigen 2 (BST2) is a single-pass type II membrane protein that belongs to the tetherin family. BST2 is predominantly expressed in the liver; lung; heart and placenta. BST2 is involved in the sorting of secreted proteins. BST2 is a human cellular protein which inhibits retrovirus infection by preventing the diffusion of virus particles after budding from infected cells. BST2 is initially discovered as an inhibitor to HIV-1 infection in the absence of Vpu; it has also been shown to inhibit the release of other viruses such as retroviruses; filoviruses; arenaviruses; and herpes viruses. BST2 may play a role in B-cell activation in rheumatoid arthritis.

For Research Use Only