Recombinant Human USP14 Protein (His Tag)

Catalog Number: PKSH033174



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-					
- 1	00	cri	m	17	٦m
J			174	, T. U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

 Species
 Human

 Mol_Mass
 48.5 kDa

 Accession
 P54578

Bio-activity Not validated for activity

Properties

Purity > 85 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, 20% Glycerol,

pH 8.0.

Reconstitution Not Applicable

Data



> 85 % as determined by reducing SDS-PAGE.

Background

Ubiquitin Carboxyl-Terminal Hydrolase 14 (USP14) belongs to the ubiquitin-specific processing (USP) family which is a deubiquitinating enzyme (DUB) with His and Cys domains. USP14 located in the cytoplasm is a proteasome-associated deubiquitinase which releases ubiquitin from the proteasome targeted ubiquitinated proteins. USP14 acts also as a physiological inhibitor of endoplasmic reticulum-associated degradation (ERAD) under the non-stressed condition by inhibiting the degradation of unfolded endoplasmic reticulum proteins via interaction with ERN1. In addition, USP14 is indispensable for synaptic development and function at neuromuscular junctions, required for the degradation of the chemokine receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis.

For Research Use Only