A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD227 Antibody[AR20.5]

catalog number: AN007620P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD227 protein

Host Mouse

Isotype Mouse IgG1, κ

Clone AR20.5


Purification >98%, Protein A/G purified

Buffer PBS, pH 7.2. Contains 0.05% Proclin300.

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



MCF-7 were stained with 0.2 µg Purified Anti-Human

CD227 Antibody[AR20.5] (Right) and 0.2 µg Mouse IgG1, κ

Isotype Control (Left), followed by Alexa Fluor® 647-

conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

MUC-1 (Mucin-1; also PEM, PEMT, Episialin, tumor-associated mucin, and carcinoma associated mucin) is a 240-450 kDa type I transmembrane glycoprotein. Human MUC-1 is 1255 amino acids (aa) in length and contains a 23 aa signal sequence, a 1135 aa extracellular domain (ECD), a 23 aa transmembrane segment, and a 74 aa cytoplasmic tail. The ECD consists of degenerate tandem repeats and a tandem repeat region, which makes up the major portion of the protein. Potential O-glycosylation sites (serines and threonines) make up more than one-fourth of the amino acids. Splicing variants produce ten isoforms for human MUC-1. Human MUC-1 is 28% aa identical to Mouse MUC-1.