

Purified Anti-Human CD227 Antibody[AR20.5]

catalog number: AN007620P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human CD227 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	AR20.5
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 µg/mL (0.5×10 ⁶ -1×10 ⁶ cells)
------------	--

Data



MCF-7 were stained with 0.2 µg Purified Anti-Human CD227 Antibody[AR20.5] (Right) and 0.2 µg Mouse IgG1, κ

Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

MUC-1 (Mucin-1; also PEM, PEMT, Episialin, tumor-associated mucin, and carcinoma associated mucin) is a 240-450 kDa type I transmembrane glycoprotein. Human MUC-1 is 1255 amino acids (aa) in length and contains a 23 aa signal sequence, a 1135 aa extracellular domain (ECD), a 23 aa transmembrane segment, and a 74 aa cytoplasmic tail. The ECD consists of degenerate tandem repeats and a tandem repeat region, which makes up the major portion of the protein. Potential O-glycosylation sites (serines and threonines) make up more than one-fourth of the amino acids. Splicing variants produce ten isoforms for human MUC-1. Human MUC-1 is 28% aa identical to Mouse MUC-1.

For Research Use Only