

## RNF144A Polyclonal Antibody

catalog number: E-AB-91827

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |                                                                                    |
|---------------------|------------------------------------------------------------------------------------|
| <b>Reactivity</b>   | Human;Rat                                                                          |
| <b>Immunogen</b>    | Recombinant fusion protein of human RNF144A                                        |
| <b>Host</b>         | Rabbit                                                                             |
| <b>Isotype</b>      | Rabbit IgG                                                                         |
| <b>Purification</b> | Affinity purification                                                              |
| <b>Buffer</b>       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

### Applications

### Recommended Dilution

|           |              |
|-----------|--------------|
| <b>WB</b> | 1:500-1:2000 |
|-----------|--------------|

### Data



Western blot analysis of extracts of various cell lines using RNF144A Polyclonal Antibody at 1:1000 dilution.

**Observed-MW:38 kDa**

### Preparation & Storage

|                 |                                                                                                          |
|-----------------|----------------------------------------------------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                                          |
| <b>Shipping</b> | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

### Background

This gene encodes a member of a family of RING finger domain-containing E3 ubiquitin ligases that also includes parkin and parc. The expression of this gene is induced by DNA damage. The encoded protein interacts with the cytoplasmic DNA-dependent protein kinase, catalytic subunit (DNA-PKcs) and promotes its degradation through ubiquitination. The orthologous mouse protein has been shown to interact with a ubiquitin-conjugating enzyme involved in embryonic development. [provided by RefSeq, Mar 2017]

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.8