# CK-19 Polyclonal Antibody

**catalog number:** **E-AB-10120**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human;Mouse;Rat |
| **Immunogen** | Recombinant protein of human KRT19 |
| **Host** | Rabbit |
| **Isotype** | Rabbit IgG |
| **Purification** | Affinity purification |
| **Buffer** | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|---|---|
| **WB** | 1:500-1:2000 |
| **IHC** | 1:100-1:300 |

## Data




Western Blot analysis of A431, Hela and Lovo cell, Human fetal lung and adult liver tissue using CK-19 Polyclonal Antibody at dilution of 1:1000

**Calculated-MW:44 kDa**



Immunohistochemistry of paraffin-embedded Rat colon using CK-19 Polyclonal Antibody at dilution of 1:50

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| **Shipping** | The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended. |

## Background

The protein encoded by this gene is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. The type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. Unlike its related family members, this smallest known acidic cytokeratin is not paired with a basic cytokeratin in epithelial cell s. It is specifically expressed in the periderm, the transiently superficial layer that envelopes the developing epidermis. The type I cytokeratins are clustered in a region of chromosome 17q12-q21.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com