


PKC zeta Polyclonal Antibody

catalog number: E-AB-92669

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human PKC zeta
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:1000
IF	1:50-1:200


Data



Western blot analysis of extracts of various cell lines using PKC zeta Polyclonal Antibody at 1:1000 dilution.


Observed-MW:70 kDa

Calculated-MW:46 kDa/56 kDa/67 kDa



Immunofluorescence analysis of A-549 cells using PKC zeta Polyclonal Antibody at dilution of 1:100 (40x lens).


Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines using PKC zeta Polyclonal Antibody at 1:1000 dilution.

Observed-MW:70 kDa

Calculated-MW:46 kDa/56 kDa/67 kDa



Immunofluorescence analysis of NIH/3T3 cells using PKC zeta Polyclonal Antibody at dilution of 1:100 (40x lens).

Blue: DAPI for nuclear staining.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8

Protein kinase C (PKC) zeta is a member of the PKC family of serine/threonine kinases which are involved in a variety of cellular processes such as proliferation, differentiation and secretion. Unlike the classical PKC isoenzymes which are calcium-dependent, PKC zeta exhibits a kinase activity which is independent of calcium and diacylglycerol but not of phosphatidylserine. Furthermore, it is insensitive to typical PKC inhibitors and cannot be activated by phorbol ester. Unlike the classical PKC isoenzymes, it has only a single zinc finger module. These structural and biochemical properties indicate that the zeta subspecies is related to, but distinct from other isoenzymes of PKC. Alternative splicing results in multiple transcript variants encoding different isoforms.