A Reliable Research Partner in Life Science and Medicine

# **RPL13A Polyclonal Antibody**

catalog number: E-AB-18938

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat

**Immunogen** Fusion protein of human RPL13A

Host Rabbit Isotype IgG


**Purification** Antigen affinity purification

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


### **Applications** Recommended Dilution

WB 1:1000-1:5000 **IHC** 1:50-1:300

### Data



Western blot analysis of LOVO Hela A549 and HT29 cell lysates using RPL13A Polyclonal Antibody at dilution of 1:1600



Immunohistochemistry of paraffin-embedded Human tonsil tissue using RPL13A Polyclonal Antibody at dilution of 1:95(×200)

# Observed-MW:Refer to figures Calculated-MW:24 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using RPL13A Polyclonal Antibody at dilution of  $1:95(\times 200)$ 

### Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

# Background

### For Research Use Only

Rev. V1.7

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a member of the L13P family of ribosomal proteins that is a component of the 60S subunit. The encoded protein also plays a role in the repression of inflammatory genes as a component of the IFN-gamma-activated inhibitor of translation (GAIT) complex. This gene is co-transcribed with the small nucleolar RNA genes U32, U33, U34, and U35, which are located in the second, fourth, fifth, and sixth introns, respectively. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed throughout the genome. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017