

## Recombinant Human AIF1 Protein (His Tag)

**Catalog Number:** PDEH100666

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human AIF1 protein Ser2-Pro147, with an C-terminal His
<b>Calculated MW</b>	17.7 kDa
<b>Observed MW</b>	16 kDa
<b>Accession</b>	P55008
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

Allograft Inflammatory Factor 1 (AIF1) contains two EF-hand domains and exists as a homodimer. AIF1 can be detected in T-lymphocytes and peripheral blood mononuclear cells. AIF1 functions as actin-binding protein that enhances membrane ruffling and RAC activation and can enhance the actin-bundling activity of LCP1. In addition, AIF1 plays a role in RAC signaling and in phagocytosis and may also in macrophage activation and function. AIF1 promotes the proliferation of vascular smooth muscle cells and of T-lymphocytes and plays a role in vascular inflammation.

### For Research Use Only