

## RAGE/AGER Monoclonal Antibody(Detector)

**catalog number: AN002030P**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                     |  |
|---------------------|--|
| <b>Reactivity</b>   | Human  |
| <b>Immunogen</b>    | Recombinant Human RAGE/AGER protein expressed by Mammalian         |
| <b>Host</b>         | Rat  |
| <b>Isotype</b>      | Rat IgG1   |
| <b>Clone</b>        | 9C7-1  |
| <b>Purification</b> | Protein A/G Purification   |
| <b>Buffer</b>       | Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300. |

### Applications Recommended Dilution

|                       |               |
|-----------------------|---------------|
| <b>ELISA Detector</b> | 0.1-0.4 µg/mL |
|-----------------------|---------------|

### Data



Sandwich ELISA-Recombinant Human RAGE/AGER protein standard curve. Background subtracted standard curve using RAGE/AGER antibody(AN002020P) (Capture), RAGE/AGER antibody(AN002030P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human RAGE/AGER protein is 78.13-5000 pg/mL.

### Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.       |
| <b>Shipping</b> | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

### Background

The advanced glycosylation end product (AGE) receptor encoded by this gene is a member of the immunoglobulin superfamily of cell surface receptors. It is a multiligand receptor, and besides AGE, interacts with other molecules implicated in homeostasis, development, and inflammation, and certain diseases, such as diabetes and Alzheimer's disease. Many alternatively spliced transcript variants encoding different isoforms, as well as non-protein-coding variants, have been described for this gene.

### For Research Use Only