

A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CD200 Protein (His Tag)

Catalog Number: PDMH100419

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Species Human

Source HEK293 Cells-derived Human CD200 protein Met1-Gly232, with an C-terminal His

 Calculated MW
 25.4 kDa

 Observed MW
 35 kDa

 Accession
 P41217

Bio-activity Not validated for activity

#### **Properties**

**Purity** > 95% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

. . . .

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

#### Data



SDS-PAGE analysis of Human CD200 proteins, 2 µg/lane of Recombinant Human CD200 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 35 kDa.

### **Background**

CD200 is a transmembrane immunoregulatory protein that belongs to the immunoglobulin superfamily. It contains one Ig like V type domain and one Ig like C2 type domain in its extracelluar domain. CD200 is widely but not ubiquitously expressed. Its receptor (CD200R) is restricted primarily to mast cells, basophils, macrophages, and dendritic cells, which suggests myeloid cell regulation as the major function of CD200. CD200 and CD200R associate via their respective N-terminal Ig-like domains. In myeloid cells, CD200R initiates inhibitory signals following receptor-ligand contact. In T cells, CD200 functions as a co-stimulatory molecule independent of the CD28 pathway. In addition, CD200 also plays an important role in prevention of graft rejection, autoimmune diseases and spontaneous abortion.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com