

A Reliable Research Partner in Life Science and Medicine

Recombinant Swine CCL2 protein(His Tag)

Catalog Number: PKSS000018

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Porcine

Source E.coli-derived Porcine CCL2 protein Gln 24-Pro 99, with an N-terminal His

Calculated MW9.4 kDaObserved MW11-17 kDaAccessionP42831

Bio-activity Not validated for activity

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin $< 0.1 \text{ EU per } \mu\text{g}$ of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

The chemokine (C-C motif) ligand 2 (CCL2), also known as monocyte chemoattractant protein (MCP)-1 and small inducible cytokine A2 (SCYA2)), is a small cytokine that belongs to the CC chemokine family responsible for monocyte attraction. Its cognate receptor, CCR2, play a critical role in regulating nociceptive processes during neuropathic pain. Both CCL2 and CCR2 are implicated in induction of autoimmunity. CCL2 recruits monocytes, memory T cells, and dendritic cells to the sites of inflammation produced by either tissue injury or infection. Recently research also showed that CCL2 might be useful as a biomarker of fibrosis as well as a target for therapeutic intervention.

For Research Use Only