

PE Anti-Mouse CD226 Antibody[10E5]

Catalog Number: E-AB-F1402D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2b, κ
Clone No.	10E5
Isotype Control	PE Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


Applications

FCM

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with APC Anti-Mouse CD8a Antibody and PE Anti-Mouse CD226 Antibody[10E5] (left) or PE Rat IgG2b, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	DNAM-1;LFA-1 associated Molecule PTA-1;PTA1 (platelet and T cell activation antigen 1);TLISA1
Uniprot ID	Q8K4F0

For Research Use Only

Gene ID

225825

Background

CD226 (DNAM-1) is constitutively expressed on native CD8+ cells and on CD4+ T cells, macrophages and NK cells. This antibody (10E5) was reported to bind about 40% of inactivated CD4+ cells and binds only to differentiated Th1 cells, but not to Th2 or Th0 cells. It is also reported to suppress antigen-specific T cell expansion and EAE (experimental allergic encephalitis) mediated by Th1 cells.