

## Recombinant Mouse IFNGR2 Protein (His Tag)

**Catalog Number:** PKSM040555

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse IFNGR2 protein Met 1-Val 243, with an C-terminal His
<b>Calculated MW</b>	26.7 kDa
<b>Observed MW</b>	40-45 kDa
<b>Accession</b>	NP_032364.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Interferon gamma receptor beta chain (IFNgammaR2), also known as IFNGR2, belongs to the type II cytokine receptor family, whose deficiency is a cause of autosomal recessive mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. This accessory factor is an integral part of the IFN-gamma signal transduction pathway and is likely to interact with GAF, JAK1, and/or JAK2. IFNGR2 is a component of the IFNgamma receptor complex along with the IFNgammaR alpha chain (IFNGR1), and is a new Bax suppressor. The C-terminal fragment (cytoplasmic domain) of IFNgammaR2 is expressed in human cancer cell lines of megakaryocytic cancer (DAMI), breast cancer (MDA-MD-468), and prostate cancer (PC3 cells). The Th1 cytokine IFNgamma, acting through its heterodimeric receptors, IFNgammaR1 and IFNgammaR2, in the induction/proliferation of Th1 cells, might suppress the Th2 responses that may underlie atopic asthma. IFNGR2 has always been seen as a key mechanism for shielding T lymphocytes from the antiproliferative effects of the IFNgamma-signal transducer and activator of transcription 1 (STAT1) pathway.