

## Biotin Anti-Mouse CD119 Antibody[GR-20]

**Catalog Number:** E-AB-F1115B

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a, κ
<b>Clone No.</b>	GR-20
<b>Isotype Control</b>	Biotin Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833B]
<b>Conjugation</b>	Biotin
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 1.0 µg per 10 <sup>6</sup> cells in 100 µL volume or 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	--

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	CD119;IFN-gamma-R-alpha;IFN-gamma-R1;Ifngr1;Interferon gamma receptor 1
<b>Uniprot ID</b>	P15261
<b>Gene ID</b>	15979
<b>Background</b>	CDw119 is a 90 kD immunoglobulin superfamily member, also known as IFN-γRα chain. It is a class II cytokine receptor family member that serves as a IFN-γ-binding chain associated with the IFN-γβ chain also known as AF-1. In addition to ligand binding, CDw119 participates in ligand trafficking. CDw119 is expressed on T and B cells, NK cells, fibroblasts, endothelial, and epithelial cells. Binding of IFN-γ induces receptor dimerization, internalization, Jak1 and Jak2 protein kinase activation and, ultimately, STAT1 activation. IFN-γ initiates and regulates a variety of immune responses.

### For Research Use Only