CALCOCO1 Polyclonal Antibody

catalog number: E-AB-52770



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Reactivity	Human;Mouse;Rat		
Immunogen	Fusion protein of human CALCOCO1		
Host	Rabbit		
Isotype	IgG		
Purification	Antigen affinity purificatio	n	
Conjugation	Unconjugated		
buffer	Phosphate buffered solution	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution		
IHC	1:30-1:150		
Data			
•	f paraffin-embedded Human cervical	Immunohistochemistry of paraffin-embedded Human	
cancer tissue using CALCOCO1 Polyclonal Antibody at		esophagus cancer tissue using CALCOCO1 Polyclonal	
Preparation & Storage	on of 1:40(×200)	Antibody at dilution of 1:40(×200)	
	Store at 2000 Valid for 12	menths Arreid frage (them arelas	
Storage		Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping		The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.	
Background			
	nding and coiled-coil domain-contain	ing protein 1), also known as cocoa, calphoglin, sarcoma	
antigen NY-SAR-3 or co	niled-coil coactivator protein is a 691	amino acid protein that shuttles between the cytoplasman	

antigen NY-SAR-3 or coiled-coil coactivator protein, is a 691 amino acid protein 1), also known as cocoa, calphogini, saconia nucleus and functions as coactivator for aryl hydrocarbon and nuclear receptors. A member of the CALCOCO family, CALCOCO1 is forms a calphoglin complex with PPA1 and PGM 1 and contains multiple functional domains through which it acts as a component of both the androgen signaling pathway and the Wnt/ β -catenin signaling pathway. CALCOCO1 exists as three alternatively spliced isoforms (termed Q9P1Z2-1, 2 and 3), which are encoded by genes mapping to human chromosome 12q13.13 and mouse chromosome 15 F3.

For Research Use Only