A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse Hemopexin/HPX Protein (His Tag)

Catalog Number: PKSM041334

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Mouse

Source HEK293 Cells-derived Mouse Hemopexin/HPX protein Ser24-Gln460, with an C-

terminal His

Calculated MW 49.8 kDa Observed MW 65-90 kDa Accession Q91X72

Bio-activity Not validated for activity

Properties

> 95 % as determined by reducing SDS-PAGE. **Purity**

Concentration Subject to label value.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.


Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles. Storage

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Supplied as a 0.2 µm filtered solution of 20mM MES, 150mM NaCl, pH 5.5. Formulation

Data



Background

Hemopexin (HPX) is plasma glycoprotein belongs to the family of the acute-phase proteins whose synthesis is induced after an inflammatory event. Hemopexin with two four-bladed beta -propeller folds has been found in other proteins including collagenases and provides sites for protein-protein interactions. The liver is the major synthesizing organ. Hemopexin participates in maintaining and recycling the iron pool by utilizing its high binding affinity toward heme composed of protoporphyrin IX and iron. It also functions in preventing oxidation caused by heme after hemolysis. Hydrophobic heme molecules can intercalate into lipid membranes and participate in the oxidation of lipid membrane components through the Fenton reaction resulting in lipid peroxidation. Hemopexin undergoes a conformational change upon the binding of heme. The conformational change allows hemopexin to interact with a specific receptor, forming a complex which is then internalized. Heme concentrations in plasma increase after hemolysis, which is associated with several pathological conditions such as reperfusion injury and ischemia.

For Research Use Only