A Reliable Research Partner in Life Science and Medicine

IDE Monoclonal Antibody

catalog number: E-AB-22041

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Synthetic Peptide

HostMouseIsotypeIgGClone1H4

Purification Protein A purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein

protectant and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000 IHC 1:50-300 IF 1:100-1:300

Data




Western Blot analysis of 1) Hela, 2) HepG2 cells using IDE Monoclonal Antibody at dilution of 1:2000.

ANGERING CONTOL

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using IDE Monoclonal Antibody at dilution of 1:200.

Observed-MW:118 kDa



Immunofluorescence analysis of Human breast tissue using

IDE Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Fax: 1-832-243-6017

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

This gene encodes a zinc metallopeptidase that degrades intracellular insulin, and thereby terminates insulins activity, as well as participating in intercellular peptide signalling by degrading diverse peptides such as glucagon, amylin, bradykinin, and kallidin. The preferential affinity of this enzyme for insulin results in insulin-mediated inhibition of the degradation of other peptides such as beta-amyloid. Deficiencies in this protein's function are associated with Alzheime r's disease and type 2 diabetes mellitus but mutations in this gene have not been shown to be causitive for these diseases. This protein localizes primarily to the cytoplasm but in some cell types localizes to the extracellular space, cell membrane, peroxisome, and mitochondrion. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described but have not been experimentally verified.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

Fax: 1-832-243-6017