

# BAX Polyclonal Antibody

catalog number: D-AB-10386L

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human;Mouse;Rat
<b>Immunogen</b>	Recombinant Human BAX Protein expressed by E.coli
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Antigen Affinity Purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4

## Applications

## Recommended Dilution

<b>WB</b>	1:500-1:1000
-----------	--------------

## Data



Western blot with BAX Polyclonal antibody at dilution of 1:1000.lane 1:MCF-7 whole cell lysate,lane 2:A549 whole cell lysate,lane 3:HepG2 whole cell lysate,lane 4:SH-SY5Y whole cell lysate,lane 5:C2C12 whole cell lysate,lane 6:NIH-3T3 whole cell lysate,lane 7:Raw264.7 whole cell lysate,lane 8:C6 whole cell lysate

**Observed-MV:20 kDa**

**Calculated-MV:4 kDa/12 kDa/15 kDa/18 kDa/19 kDa/21 kDa/24 kDa**

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

## Background

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2,and functions as an apoptotic activator. This protein is reported to interact with,and increase the opening of,the mitochondrial voltage-dependent anion channel (VDAC),which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants,which encode different isoforms,have been reported for this gene.

## For Research Use Only