Recombinant Human NPRC/NPR3 (C-Fc)

Catalog Number: PKSH034022



Description	
Species	Human
Mol_Mass	77.5 kDa
Accession	P17342
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

> 95 % as determined by reducing SDS-PAGE.

kDa

120 90

60

40

30

MK

R

Background

Atrial Natriuretic Peptide Receptor-3 (NPR3), also known as NPRC or ANPR-C, is one of the three natriuretic peptide receptors, is a type I transmembrane glycoprotein. The natriuretic system is key to the maintenance of vascular tone and cardiovascular homeostasis. Receptor for the natriuretic peptide hormones, binding with similar affinities atrial natriuretic peptide NPPA/ANP, brain natriuretic peptide NPPB/BNP, and C-type natriuretic peptide NPPC/CNP. May function as a clearance receptor for NPPA, NPPB and NPPC, regulating their local concentrations and effects. Osteocrin was found to be a specific ligand to NPR3. NPR3 is necessary for Osteocrin to regulate femoral, tibial, and metatarsal bone elongation.

For Research Use Only