

## Recombinant Human Decorin Protein (Human Cells, His Tag)

Catalog Number: PKSH033700

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Decorin protein Gly17-Lys359, with an C-terminal His
<b>Calculated MW</b>	38.8 kDa
<b>Observed MW</b>	43-50 kDa
<b>Accession</b>	P07585
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Decorin is a secreted chondroitin/dermatan sulfate proteoglycan in the family of small leucine-rich proteoglycans (SLRPs). SLRP family members are characterized by N-terminal and C-terminal cysteine-rich regions which flank the central region containing 10-12 tandem leucine-rich repeats (LRR). The human Decorin cDNA encodes a 359 amino acid (aa) precursor that includes a 16 aa signal sequence and a 14 aa propeptide. Alternate splicing of human Decorin generates five isoforms with variable length deletions. Decorin is an N-glycosylated protein that also carries a variably sized hybrid chondroitin/dermatan sulfate chain at Ser34. Decorin regulates assembly of the extracellular collagen matrix and the bioactivity of the matrix associated growth factors FGF2; GDF8/Myostatin; TGFβ; and WISP1. It also binds and activates EGF R; ErbB4; and IGFI-R. In vivo; Decorin promotes myoblast differentiation; supports angiogenesis; and inhibits tumor progression.