Elabscience®

Immunohistochemistry of paraffin-embedded mouse heart

using COX7A1 Polyclonal Antibody at dilution of 1:100

(40x lens).Perform high pressure antigen retrieval with 10

mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

COX7A1 Polyclonal Antibody

catalog number: E-AB-90936

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human COX7A1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200

Data



Western blot analysis of extracts of Mouse heart using COX7A1 Polyclonal Antibody at 1:500 dilution. **Observed-MW:9 kDa**

Calculated-MW:9 kDa



Immunohistochemistry of paraffin-embedded rat heart using COX7A1 Polyclonal Antibody at dilution of 1:100 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining

 Preparation & Storage

 Storage
 Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

 Shipping
 The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

 Background
 Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 1 (muscle isoform) of subunit VIIa and the polypeptide 1 is present only in muscle tissues. Other polypeptides of subunit VIIa are present in both muscle and nonmuscle tissues, and are encoded by different genes.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com