

FITC Anti-Human CD151 Antibody[50-6]

Catalog Number: AN00635C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	50-6
Isotype Control	FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C]
Conjugation	FITC
Conjugation Information	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood platelets are stained with FITC Anti-Human CD151 Antibody[50-6] (filled gray histogram) or FITC Mouse IgG1, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	Platelet-endothelial tetraspan antigen 3;GP27;Membrane glycoprotein SFA-1; Tetraspanin-24;Tspan-24;CD151
Uniprot ID	P48509
Gene ID	977

For Research Use Only

Background

CD151, also known as PETA-3, is a 29 kD protein member of the tetraspanin family (TM4SF). As other members in the family, PETA-3 is composed of two extracellular loops (one larger than the other), four transmembrane domains, and two cytoplasmic tails. Also, as other members in the family, CD151 is involved in cell adhesion, migration, and tumor cell metastasis. PETA-3 is expressed by many different cell types such as immature hematopoietic cells, megakaryocytes, platelets, keratinocytes, epithelial cells, muscle cells, Schwann cells, and vascular endothelium. On the cell surface, PETA-3 associates with CD9, CD181, and the integrins $\alpha 3\beta 1$, $\alpha 5\beta 1$, and $\alpha 6\beta 4$.