Elabscience®


EEF1G Polyclonal Antibody


catalog number: E-AB-65564

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human EEF1G (NP_001395.1).
Host	Rabbit
Is otype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:100
IF	1:50-1:100

Data






EEF1G Polyclonal Antibody at dilution of 1:100 (40x lens).


Western blot analysis of extracts of various cell lines using EEF1G Polyclonal Antibody at dilution of 1:1000.

Observed-MW:50 kDa Calculated-MW:50 kDa/56 kDa




Immunohistochemistry of paraffin-embedded Human esophagus using EEF1G Polyclonal Antibody at dilution of

1:100 (40x lens).



Immunofluorescence analysis of C6 cells using EEF1G Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Elabscience®

Immunofluorescence analysis of NIH-3T3 cells using EEF1G Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.

Background

This gene encodes a subunit of the elongation factor-1 complex, which is responsible for the enzymatic delivery of aminoacyl tRNAs to the ribosome. This subunit contains an N-terminal glutathione transferase domain, which may be involved in regulating the assembly of multisubunit complexes containing this elongation factor and aminoacyl-tRNA synthetases.

For Research Use Only