

A Reliable Research Partner in Life Science and Medicine

VCAM-1/CD106 Polyclonal Antibody(Capture/Detector)

catalog number: AN000250P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

r .	
LIACCELL	ntion
Descri	JULUI

Reactivity Human; Mouse; Rat

Immunogen Recombinant Rat VCAM-1/CD106 protein expressed by Mammalian

Host Rabbit
Isotype Rabbit IgG

Purification Antigen Affinity Purification

Conjugation Unconjugated

Buffer Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications	Recommended Dilution
WB	1:500-1:1000
IHC	1:1000-1:2000
ELISA Capture	2-8 μg/mL
ELISA Detector	0.1 - $0.4 \mu g/mL$


Web: www.elabscience.cn

Data



Western Blot with VCAM-1/CD106 Polyclonal Antibody at dilution of 1:500.Lane 1:Mouse spleen,Lane 2:Rat kidney,Lane 3:Rat spleen.

Observed-MV:75-100 kDa Calculated-MV:81 kDa



Sandwich ELISA-Recombinant Rat VCAM-1/CD106 protein standard curve.Background subtracted standard curve using

VCAM-1/CD106 antibody(AN000250P)

(Capture), VCAM-1/CD106 antibody(AN000250P)(Detector) in sandwich ELISA. The reference range value for Recombinant Rat VCAM-1/CD106 protein is 31.25-2000

pg/mL.

Preparation & Storage

For Research Use Only

Immunohistochemistry of paraffin-embedded Human Kidney cancer using VCAM-1/CD106 Polyclonal Antibody at dilution of 1:2000.



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Cell adhesion glycoprotein predominantly expressed on the surface of endothelial cells that plays an important role in immune surveillance and inflammation. Acts as a major regulator of leukocyte adhesion to the endothelium through interaction with different types of integrins. During inflammatory responses, binds ligands on the surface of activated endothelial cells to initiate the activation of calcium channels and the plasma membrane-associated small GTPase RAC1 leading to leukocyte transendothelial migration. Serves also as a quality-control checkpoint for entry into bone marrow by providing a 'don't-eat-me' stamping in the context of major histocompatibility complex (MHC) class-I presentation.