

A Reliable Research Partner in Life Science and Medicine

# FITC Anti-Mouse Foxp3 Antibody[3G3]

Catalog Number: E-AB-F1238C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse Mouse Host

Mouse IgG1, κ Isotype

Clone No. 3G3

FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C] Isotype Control

Conjugation

**Conjugation Information** FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

#### **Applications** Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



C57BL/6 splenocytes are surface stained with APC Anti-Mouse CD4 Monoclonal Antibody followed by fixation and permeabilization using the Foxp3 Staining/Transcription Factor Buffer Set and intracellular staining with FITC Anti-Mouse Foxp3 Monoclonal Antibody.

### **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

# **Antigen Information**

**Alternate Names** Forkhead box protein P3;Foxp3;IPEX;JM2

**Uniprot ID** 

Gene ID 50943;20371;317382

# For Research Use Only

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com

#### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

## **Background**

FOXP3 is a 50-55 kD transcription factor, also known as Forkhead box protein P3, Scurfin, JM2, or IPEX. It is proposed to be a master regulatory gene and more specific marker of T regulatory cells than most cell surface markers (such as CD4 and CD25). Transduced expression of FOXP3 in CD4+/CD25- cells has been shown to induce GITR, CD103, and CTLA4 and impart a T regulatory cell phenotype. FOXP3 is mutated in X-linked autoimmunity-allergic dysregulation syndrome (XLAAD or IPEX) in humans and in 'scurfy' mice. Overexpression of FOXP3 has been shown to lead to a hypoactive immune state suggesting that this transcriptional factor is a central regulator of T cell activity. In human, unlike in mouse, two isoforms of FOXP3 have been reported: one ( FOXP3) corresponding to the canonical full-length sequence; the other (FOXP3  $\delta 2$ ) lacking exon 2.

Fax: 1-832-243-6017