

## Recombinant ABHD14B Monoclonal Antibody

catalog number: **AN300446P**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*


### Description

<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human ABHD14B Protein
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Clone</b>	2D7
<b>Purification</b>	Protein A
<b>Buffer</b>	0.2 µm filtered solution in PBS

### Applications Recommended Dilution

<b>ICC/IF</b>	1:20-1:100
---------------	------------

### Data



Immunofluorescence analysis of ABHD14B in Hela cells.

Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human ABHD14B monoclonal antibody (1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor® 594-conjugated Goat Anti-rabbit IgG secondary antibody (red) and counterstained with DAPI (blue). Positive staining was localized to nucleus.

### Preparation & Storage

<b>Storage</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Shipping</b>	Ice bag

### Background

ABHD14B belongs to the AB hydrolase superfamily, ABHD14 family. It can be detected in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, placenta, lung, liver, skeletal muscle, pancreas and kidney. ABHD14B has hydrolase activity towards p-nitrophenyl butyrate (in vitro) and may interact with TAF1. It may activate transcription. Recombinant human ABHD14B protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques. ABHD14B contains an alpha/beta hydrolase fold, which is a catalytic domain found in a very wide range of enzymes. In molecular biology, the alpha/beta hydrolase fold is common to a number of hydrolytic enzymes of widely differing phylogenetic origin and catalytic function. The Ab hydrolase domain containing gene subfamily is comprised of 15 mostly uncharacterized members.

### For Research Use Only