

## ASAH2 Polyclonal Antibody

catalog number: E-AB-14747

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human;Mouse;Rat
<b>Immunogen</b>	Recombinant protein of human ASAH2
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Conjugation</b>	Unconjugated
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications

### Recommended Dilution


<b>WB</b>	1:200-1:1000
<b>IHC</b>	1:25-1:100

### Data




Western Blot analysis of Lovo cell and Mouse heart tissue using ASAH2 Polyclonal Antibody at dilution of 1:270

**Calculated-MW:86 kDa**



Immunohistochemistry of paraffin-embedded Human colon cancer using ASAH2 Polyclonal Antibody at dilution of 1:50



Immunohistochemistry of paraffin-embedded Human brain using ASAH2 Polyclonal Antibody at dilution of 1:50

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

### Background

Ceramidases (EC 3.5.1.23), such as ASAH2, catalyze hydrolysis of the N-acyl linkage of ceramide, a second messenger in a variety of cellular events, to produce sphingosine. Sphingosine exerts both mitogenic and apoptosis-inducing activities, and its phosphorylated form functions as an intra- and intercellular second messenger.

### For Research Use Only