

AHR Polyclonal Antibody

catalog number: **E-AB-62004**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse
Immunogen	Recombinant fusion protein of human AHR (NP_001612.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications


Recommended Dilution

IHC	1:50-1:200
------------	------------

Data



Immunohistochemistry of paraffin-embedded Human stomach using AHR Polyclonal Antibody at dilution of 1:100 (40x lens).



Western blot analysis of various lysates using AHR Rabbit pAb (A1451SP) at 1:10000 dilution incubated overnight at 4°C. [Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. |Lysates/proteins: 25 µg per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit (RM00020). |Negative control (NC): Jurkat. |Exposure time: 3 s.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The protein encoded by this gene is a ligand-activated helix-loop-helix transcription factor involved in the regulation of biological responses to planar aromatic hydrocarbons. This receptor has been shown to regulate xenobiotic-metabolizing enzymes such as cytochrome P450. Before ligand binding, the encoded protein is sequestered in the cytoplasm; upon ligand binding, this protein moves to the nucleus and stimulates transcription of target genes.

For Research Use Only