A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Mouse CD183/CXCR3 Antibody[CXCR3-173]

Catalog Number: GF11140

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

**Reactivity** Mouse

Immunogen Recombinant Mouse CD183/CXCR3 protein

Host Armenian Hamster Isotype Armenian Hamster IgG

Clone CXCR3-173

**Purification** >98%, Protein A/G purified

Conjugation None (AF/LE)


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the

LAL method.

# Applications Recommended Dilution

FCM  $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$ 

#### **Data**



C57/BL6 Mouse splenocytes were stained with 0.2 μg AF/LE Purified Anti-Mouse CD183/CXCR3 Antibody[CXCR3-173] (Right) and 0.2 μg Armenian Hamster IgG, κ Isotype Control (Left), followed by PE-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody.

## **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid

freeze / thaw cycles. This preparation contains no preservatives, thus it

should be handled under as eptic conditions.

Shipping Ice bag

# **Background**

CD183/CXCR3, also known as CXCR3, is a member of the C-X-C chemokine family, characterized by a pair of cysteine residues separated by a single amino acid. CXCR3 is a 38 kD seven pass transmembrane receptor coupled to G-protein. It mediates Ca2+ mobilization and chemotaxis in response to C-X-C chemokines, such as IP10 (CXCL10), MIG (CXCL9), I-TAC (CXCL11) and PF4 (CXCL4). CXCR3 is expressed primarily on activiated T lymphocytes, NK cells, and some epithelial cells and endothelial cells. It is not expressed on B cells, monocytes or granulocytes.