Recombinant Human GM-CSF Protein(His Tag)

Catalog Number: PDMH100207



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						٠,		
D	Α		C	PТ	m	ŤΤ	U.	m
\mathbf{L}	v	0	v.	ш	עו	UΙ	U.	

Species Human

Source Mammalian-derived Human GM-CSF proteins AIa18-Glu144, with an C-terminal His

 Mol_Mass
 13.9 kDa

 Accession
 P04141

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human GM-CSF proteins, 2 μ g/lane of Recombinant Human GM-CSF proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 13.9 KD

Background

CSF2 (Colony Stimulating Factor 2) is a Protein Coding gene. Diseases associated with CSF2 include Mucositis and Candida Glabrata. Among its related pathways are RET signaling and IL-15 Signaling Pathways and their Primary Biological Effects in Different Immune Cell Types. GO annotations related to this gene include cytokine activity and granulocyte macrophage colony-stimulating factor receptor binding. The protein encoded by this gene is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer. This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known to be associated with interstitial deletions in the 5q-syndrome and acute myelogenous leukemia. Other genes in the cluster include those encoding interleukins 4, 5, and 13.

For Research Use Only