

## Recombinant Human Nogo Receptor/NgR Protein (His &Fc Tag)

**Catalog Number:** PKSH031590

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Nogo Receptor/NgR protein Met 1-Ser 447, with an C-terminal His & Fc
<b>Calculated MW</b>	73.0 kDa
<b>Observed MW</b>	100 kDa
<b>Accession</b>	NP_075380.1
<b>Bio-activity</b>	Measured by its ability to bind recombinant human RTN4 in a functional ELISA.

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V3.5

Reticulon 4 receptor (RTN4R), also known as Nogo-66 Receptor (NgR), is a glycosylphosphoinositol (GPI)-anchored protein that belongs to the Nogo receptor family including three members. Mouse RTN4R cDNA contains 10 LRP (Leucine-rich) repeats. RTN4R is expressed predominantly in neurons and their axons in the central nervous systems (CNS). As a receptor for myelin-derived proteins Nogo, myelin-associated glycoprotein (MAG), and myelin oligodendrocyte glycoprotein (OMG), RTN4R mediates axonal growth inhibition and may play a role in regulating axonal regeneration and plasticity in the adult CNS. It has been shown that RTN4R performs its inhibitory actions by interacting with the p75 neurotrophin receptor (p75NTR), a TNFRSF member also known for modulating the activities of the Trk family and for inducing apoptosis in neurons and oligodendrocytes. RTN4R may be proposed as a potential drug target for treatment of various neurological conditions such as spinal cord injury, CNS lesions, peripheral nerve injury, stroke and Alzheimer's disease (AD). Additionally, RTN4R may play a role in regulating the function of the gap junctions.