



A Reliable Research Partner in Life Science and Medicine

Human CD38 Antibody Pair Set

Catalog No. E-KAB-0700 Applications ELISA

Synonyms T10;p45;cADPRH;CD38

Kit components & Storage

Title	Specifications	Storage
Human CD38 Capture Antibody	1 vial, 100 μ g	Store at -20°C. Avoid freeze / thaw
		cycles.
Human CD38 Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze / thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0700)		
		Human CD38 Capture Antibody	Human CD38 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human CD38 protein	Recombinant Human CD38 protein	
Information	Swissprot	P28907		
Product details	Reactivity	Human	Human	
	Host	Sheep	Sheep	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Affinity purification	Affinity purification	
	Specificity	Detects Human CD38 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Human CD38 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Human CD38 Capture	
Capture		Antibody	10
			Optical Density
ELISA	1:1000-1:10000	Human CD38 Detection	00 O O O O O O O O O O O O O O O O O O
Detection		Antibody (Biotin)	
			0.01 100 1000 10000 Human CD38 Concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-hematopoietic tissues (such as brain,muscle,and kidney). In humans,it is expressed at high levels on plasma cells and activated T and B cells,natural killer (NK) lymphocytes,myeloblasts,and erythroblasts. By functioning as both a cyclase and a hydrolase,CD38 mediates lymphocyte activation,adhesion,and the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com