

Recombinant Mouse ICOS/AILIM Protein (His Tag)

Catalog Number: PKSM041217

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse ICOS/AILIM protein Glu21-Leu142, with an C-terminal His
Calculated MW	14.7 kDa
Observed MW	16-30 kDa
Accession	Q9WVS0
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

Inducible Costimulator(ICOS) is a member of the growing CD28 family of immune costimulatory receptors. Other family members are CD28, CTLA4 and PD1. ICOS shares approximately 39% amino acid similarity with CD 28 and CTLA4. Mouse and human ICOS share approximately 72% amino acid identity. ICOS is expressed on most CD45RO+ cells. ICOS expression is up-regulated within approximately 24-48 hours of activation on Th primed cells. B7-H2, a member of the B7 family of costimulatory ligands, has been identified as the ICOS ligand. The B7-H2/ ICOS interaction appears to play roles in T cell dependent B cell activation and Th differentiation. In addition, ICOS is more potent in the induction of IL-10 production, acytokine important for suppressive function of T regulatory cells.

For Research Use Only