



A Reliable Research Partner in Life Science and Medicine

Porcine LDLR Antibody Pair Set

Catalog No. E-KAB-0673 Applications ELISA

Synonyms FH;FHC;LDLCQ2;Familial Hypercholesterolemia

Kit components & Storage

Title	Specifications	Storage
Porcine LDLR Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Porcine LDLR Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0673)	
		Porcine LDLR Capture Antibody	Porcine LDLR Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Porcine LDLR protein	Recombinant Porcine LDLR protein
Information	Swissprot	Q28832	
Product details	Reactivity	Porcine	Porcine
	Host	Rabbit	Rabbit
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Protein A & Antigen Affinity	Protein A & Antigen Affinity
	Specificity	Detects Porcine LDLR in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Porcine LDLR Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Porcine LDLR Capture	
Capture		Antibody	10
			8 13
			Optical Density
ELISA	1:1000-1:10000	Porcine LDLR Detection	O O O Diffe
Detection		Antibody (Biotin)	
			0.01
			Poreine LDLR Concentration (ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

LDLR (low density lipoprotein receptor) is a member of the LDL receptor gene family and is involved in receptor-mediated endocytosis of specific ligands. The LDLR is a cell surface glycoprotein that scavenges LDL from the blood and regulates plasma LDL cholesterol. The cytoplasmic domain of the LDL receptor is necessary for the receptor to cluster in coated pits; which promotes the rapid endocytosis of bound LDL. The protein is highly glycosylated through N- and O-linkages and thus migrates at 100 to 160 kDa bands on SDS-PAGE.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com