

## Recombinant FGFR3 Monoclonal Antibody

catalog number: **AN301784L**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human;
<b>Immunogen</b>	Recombinant human FGFR3 fragment
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG, $\kappa$
<b>Clone</b>	A492
<b>Purification</b>	Protein A purified
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications

### Recommended Dilution


<b>WB</b>	1:1000-1:2000
<b>IP</b>	1:25-1:50

### Data



Western Blot with FGFR3 Monoclonal Antibody at dilution of 1:2000. Lane 1: HepG2, Lane 2: HEK-293, Lane 3: K562, Lane 4: MCF-7

**Observed-MW:125 kDa**  
**Calculated-MW:88 kDa**



Immunoprecipitation analysis using anti-FGFR3 Monoclonal Antibody. Western blot was performed from the immunoprecipitate using FGFR3 Monoclonal Antibody at a dilution of 1:50. Lane 1: 10% Input, Lane 2: FGFR3 Monoclonal Antibody, Lane 3: Rabbit monoclonal IgG Isotype

**Observed-MW:125 kDa**  
**Calculated-MW:88 kDa**

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

Fibroblast growth factor receptor 3 is a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid sequence being highly conserved between members and among divergent species. The FGFR proteins are involved in a wide array of pathways known to play a significant role in cancer. Activation of these receptors can lead to activation of the RAS-MAPK pathway and the PI3K-AKT pathway, among others.

### For Research Use Only