

## Purified Anti-Human/Monkey CD16 Antibody[3G8]

catalog number: E-AB-F1236A

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


|                     |   |
|---------------------|---|
| <b>Reactivity</b>   | Human;Rhesus;Cynomolgus   |
| <b>Immunogen</b>    | Recombinant Human CD16 protein  |
| <b>Host</b>         | Mouse   |
| <b>Isotype</b>      | Mouse IgG1, κ   |
| <b>Clone</b>        | 3G8   |
| <b>Purification</b> | >98%, Protein A/G purified  |
| <b>Buffer</b>       | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.<br>Dialyze to completely remove the stabilizer prior to labeling. |

### Applications

### Recommended Dilution

|            |   |
|------------|---|
| <b>FCM</b> | 2 µg/mL(0.5×10 <sup>6</sup> -1×10 <sup>6</sup> cells) |
|------------|---|

### Data



Human peripheral blood lymphocytes were stained with 0.2 µg Purified Anti-Human CD16 Antibody[3G8] (Right) and 0.2 µg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD56 PE-conjugated Monoclonal Antibody.

### Preparation & Storage

|                 |  |
|-----------------|--|
| <b>Storage</b>  | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag  |

### Background

CD16 is a 60 kD highly glycosylated protein. It is a member of the Ig superfamily and is also known as B7-1, B7, and Ly-53. CD16 is constitutively expressed on dendritic cells and monocytes/macrophages, and inducibly expressed on activated B and T cells. The ligation of CD28 on T cells with CD16 and CD86 (B7-2) on antigen presenting cells (such as dendritic cells, macrophages, and B cells) elicits co-stimulation of T cells resulting in enhanced cell activation, proliferation, and cytokine production. CD16 appears to be expressed later in the immune response than CD86. CD16 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

### For Research Use Only