

A Reliable Research Partner in Life Science and Medicine

PE/Cyanine 5 Anti-Human CD48 Antibody [156-4H9]

Catalog Number: E-AB-F1061G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ **Clone No.** 156-4H9

Isotype Control PE/Cyanine5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792G]

Conjugation PE/Cyanine 5

Conjugation Information PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Human peripheral blood lymphocytes are stained with PE/Cyanine5 Anti-Human CD48 Antibody (filled gray histogram) or Mouse IgG1 Isotype Control PE/Cyanine5 (empty black histogram).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names BCM1 surface antigen;BLAST-1;CD48;CD48 antigen;Cd48;HM48-1;MRC OX-45

surface antigen; SLAMF2; sgp-60

Uniprot ID P09326

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 962

CD48 is a 40-47 kD GPI-anchored membrane protein, also known as Blast-1 and HuL y-m3. It is a member of the CD2 family that contains 2 IgSF domains and is widely expressed on both resting and activated hematopoietic cells with the exception of granulocytes, platelets, and erythrocytes. CD48 binds to CD2 at a considerably (>100-fold) lower affinity than CD58. It is thought to contribute to T cell activation. The cytoplasmic tail of CD48 has been shown to bind to the kinases Lck and Fyn.