

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Red 780 Anti-Rat CD4(domain 1) Antibody[OX-38]

Catalog Number: E-AB-F1105S

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Rat
Host Mouse

Isotype Mouse IgG2a, κ

Clone No. OX-38

Isotype Control Elab Fluor® Red 780 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802S]

Conjugation Elab Fluor[®] Red 780

Conjugation Information Elab Fluor[®] Red 780 is designed to be excited by the Red (627-640 nm) laser and

detected using an optical filter centered near 770 nm (e.g., a 780/60 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Rat splenocytes are stained with Elab Fluor® Red 780 Anti-Rat CD4(domain 1) Antibody[OX-38] (filled gray histogram)

or Elab Fluor[®] Red 780 Mouse IgG2a, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;domain 1

 Uniprot ID
 P05540

 Gene ID
 24932

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD4, also known as T4, is a 55kD glycoprotein member of the immunoglobin superfamily and is expressed on majority of thymocytes, macrophages, and a peripheral T cell subset (T helper cells). CD4 is a T cell co-receptor that interacts with the MHC class II molecule and is involved in T cell activation.