A Reliable Research Partner in Life Science and Medicine

Recombinant Human Contactin 2/CNTN2 Protein (His Tag)

Catalog Number: PKSH032278

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human Contactin 2;CNTN2 protein Ser31-Asn1012, with an C-

terminal His

Calculated MW 108.5 kDa
Observed MW 110-140 kDa
Accession Q02246

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Contactin-2 (CNTN2) is encoded by the CNTN2 gene, which belongs to the immunoglobulin superfamily and contactin family. It contains 4 fibronectin type-III domains and 6 Ig-like C2-type domains. It is a glycosylphosphatidylinositol (GP I)-anchored neuronal membrane protein that functions as a cell adhesion molecule. CNTN2 may play a role in the formation of axon connections in the developing nervous system. In conjunction with another transmembrane protein, CNTNAP2, contributes to the organization of axonal domains at nodes of Ranvier by maintaining voltage-gated potassium channels at the juxtaparanodal region. It may also be involved in glial tumorigenesis and may provide a potential target for therapeutic intervention.

For Research Use Only