Recombinant Mouse NBL1/DAND1 Protein (His Tag)

Catalog Number: PKSM041115

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Mouse
Source	HEK293 Cells-derived Mouse NBL1/DAND1 protein Ala17-Asp178, with an C-
	terminal His
Calculated MW	18.4 kDa
Observed MW	25-28 kDa
Accession	Q61477
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Elabscience®

Differential screening-selected gene aberrative in neuroblastoma (DAN) is a member of the DAN family of secreted glycoproteins. DAN family antagonists are characterized by a DAN domain that contains a cystine knot motif which is essential for binding to BMP ligands. Members of this family include DAN, gremlin, protein related to DAN and cerberus (PRDC), cerberus, sclerostin (SOST) and uterine sensitization-associated gene 1 protein, and control diverse processes in growth, development and the cell cycle. It has also been reported that DAN family plays crucial role in early mouse embryo development by inhibiting the action of bone morphogenic proteins and modulating the action of transforming growth factor-β superfamily members. DAN is synthesized by small-to intermediate-sized DRG neurons and transported to the sensory nerve terminals in the skin or to the sensory nerve terminals in the dorsal horn. It has been reported that DAN is ubiquitously expressed in adult rat and human tissues. Morphological studies have revealed that, in adult rat, DAN mRNA is expressed ubiquitously in lung and brain, but not in liver.