

Recombinant Mouse IL-17D protein(N-His)

Catalog Number: PKSM041506

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Mouse
Source	E.coli-derived Mouse IL-17D protein Ala 25-Arg 205, with an N-terminal His
Calculated MW	20.7 kDa
Observed MW	25 kDa
Accession	NP_665836.2
Bio-activity	Not validated for activity

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20 mM sodium citrate, 0.2 M NaCl, pH 4.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

The Interleukin-17 family proteins, comprising six members (IL-17, IL-17B through IL-17F), are secreted, structurally related proteins that share a conserved cysteine-knot fold near the C-terminus, but have considerable sequence divergence at the N-terminus. IL-17 family proteins are proinflammatory cytokines that induce local cytokine production and are involved in the regulation of immune functions. Among IL-17 family members, IL-17D is most closely related to IL-17B, sharing 27% aa sequence homology. IL-17D is expressed preferentially in skeletal muscle, heart, adipose tissue, lung, pancreas, and nervous system. Like other IL-17 family members, IL-17D modulates immune responses indirectly by stimulating the production of myeloid growth factors and chemokines including IL-6, IL-8, and GM-CSF. IL-17D has also been shown to suppress the proliferation of myeloid progenitors in colony formation assays.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017